

MONITORING

of golden-headed lion tamarins in cocoa-agroforest



This project monitors the endangered golden-headed lion tamarins (*Leontopithecus chrysomelas*) in cocoa-agroforestry systems (so called cabucas) to determine ecological pressures and to understand baseline habitat suitability. This will allow us to identify which management practices will enable the long-term survival of golden-headed lion tamarins in the wild.

AMAP is a German-Brazilian environmental organization dedicated to the conservation of the Mata Atlântica, the Brazilian Atlantic Rain forest. Especially the long term survival of golden-headed lion tamarins is in the focus of AMAP's mission, as an flagship species of the region. AMAP promotes the conservation of the Mata Atlântica through land acquisition, reforestation, the support of research projects and biodiversity-friendly cocoa cultivation. The base of our activities in the region is the association's own farm "Bom Pastor" not far from the Almada River.

Project coordinator

Dr. Joanison Vicente dos Santos Teixeira

Post Doc at the LECAP (Laboratório de Etnoconservação e Áreas Protegidas), Departamento de Ciências Agrárias e Ambientais; Universidade Estadual de Santa Cruz (UESC); Universidade Estadual de Santa Cruz (UESC) and
Project coordinator AMAP Brazil

Project coordinator and initiator of the project

Prof. Dr. Leonardo de Carvalho Oliveira

Ecology and Biodiversity Conservation, Universidade Estadual de Santa Cruz (UESC); Professor Adjunto da Universidade do Estado do Rio de Janeiro (FFP-UERJ)

Cooperation partner, funding

Christian Wolff

Coordinator for Reforestation and Biodiversity
2. Executive Director AMAP Brazil and Germany

christian.wolff@amap-brazil.org

Tel.: +49 178 3370845

The golden-headed lion tamarin is an endangered primate, endemic to a small part of Southern Bahia within the Brazilian Atlantic Forest. Caused by anthropogenic habitat reduction and fragmentation, the long-term survival of the species depends on its ability to persist in cocoa agroforests (cabruças). In the municipality of Ilhéus, our project area, predominant cabruças forms as matrix a connected forest landscape. Since 2008, several groups of lion tamarins are monitored. The monitoring generates a biological, ecological and behavioral dataset, which gives us the knowledge to understand the basic habitat suitability of cabruças and to develop biodiversity friendly cocoa harvesting and maintenance strategies for local farmers. This strategy will secure the long-time survival of golden-headed lion tamarins in a human dominated agricultural landscape.

I. Current situation

The golden-headed lion tamarin (*Leontopithecus chrysomelas*) is an endangered small primate endemic to southern Bahia in the Brazilian Atlantic Forest, the Mata Atlântica. The Mata Atlântica is one of the most threatened tropical forests and was especially during the 20th century reduced to 10% of its original extent. Nevertheless, it is one of the most diverse regions in the world (Myers et al. 2000; Shi et al. 2005). The remaining forest is strongly fragmented, within the lion tamarin range only 5% of the forest patches are larger than 36 ha (Zeigler et al. 2010). The home range size varies from 40 to 197 ha (Dietz et al. 1996; Rylands 1993) with an average of 83 ha (Oliveira et al. 2011). Thus, the remaining habitat is also extremely reduced. Estimates of the total wild population size ranges from 6,000 to 15,000 individuals in an area of ca. 19,000km² (Pinto & Rylands 1997), however, many small populations, mainly in the western part of the distribution, disappeared in the past years reducing the geographic distribution of lion tamarins (Raboy et al. 2010). Results from a Population and Habitat Viability Analysis

suggested that only one population, situated in the protected Una Biological and Wildlife Refuge, is viable and capable of preserving sufficient genetic variability for a period of 100 years (Zeigler et al. 2010). However, assuming a metapopulation scenario in which forested areas are connected by a matrix habitat, in this case cocoa agroforest (so called cabruça), suitable for dispersal of lion tamarins, their conservation in the wild is relatively secure (Holst et al. 2006).



Cabruça of Fazenda Bom Pastor, habitat of lion tamarins.

Agroforest can be defined as a complex multistrata agroforestry systems based on shade-tolerant understorey crops such as cocoa (*Theobroma cacao*) or coffee (*Coffea* spp.) grown under a complex and often species-rich canopy of native and/or planted trees (Schroth et al., 2014). This forest-agriculture-system can thus provide habitat and resources for forest-dependent species that would not survive in a purely agricultural landscape, or may permit species dispersal in a fragmented landscape (Schroth et al., 2007). The cabruça has been considered as an important habitat for the Mata Atlântica biodiversity for plants (Sambuichi 2002, 2006; Sambuichi and Haridasan 2007) and animals (Pardini 2004; Delabie et al., 2007; Faria et al., 2006 and 2007).

In contrast to many other forest dwelling animals, lion tamarins and other marmosets (Callitrichidae), uses disturbed and regenerating forests for foraging, for instance fruits, flowers, nectar, gum and animal prey (Oliveira et al., 2010). Especially fruits are available in large

quantities in secondary forests and cabruças, caused by a better light availability as in primary forests. Cabruças (Oliveira pers.comm.) provide more than 60% of the habitable areas for lion tamarins in the eastern part of its distribution. Cabruças can provide suitable conditions, moreover lion tamarins are in many areas the last remaining frugivorous mammals, putting them in a key role as seed vector for forest regeneration (Oliveira & Estrada 2017).



A golden-headed lion tamarin of the Fazenda Bom Pastor

Thus, conservation approaches are concentrated on two areas with viable lion tamarin populations, the Una Biological and Wildlife Refuge as a larger protected forest area and the cocoa region of Ilhéus. The latter one, our project area, represents a patchwork of different vegetation types embedded in a cabruça landscape matrix. Primary and secondary forest fragments are present, as well as pastures. Cocoa is the most important crop of the Ilhéus region and plays a major role in local economics and is the only income generating agricultural system, which allows the implementation of biodiversity conservation approaches.

The evaluation of the use of cabruça by lion tamarins was recognized as a conservation priority for the species (Holst et al. 2006). Our long-term research project wants to understand how different management types of cabruças shape the suitability as habitat. Therefore an analysis of biological, ecological and behavioral datasets are necessary.

Radio collared lion tamarin groups are monitored since 2008 by Dr. Leonardo Oliveira (Universidade Estadual de Santa Cruz). During

the monitoring in our project area it was possible to document that lion tamarins exclusively survive and reproduce in cabruças (Oliveira et al. 2011). Moreover, in cases where cabruça contains concentrated resources, such as jackfruit and bromeliads lion tamarins are heavier and bigger than expected for the species, with a higher reproduction (Oliveira et al., 2011). In one monitored group (the Santa Rita group) the first report of triplets in wild populations could be made, additionally this group shows the smallest home range (22-28 ha) and highest density reported (0,17 Ind./ha) for lion tamarins (Oliveira et al., 2011). A categorized list of key tree species crucial for the survival of lion tamarins in the cabruças was created by Oliveira et al. 2010.

The long-term monitoring project aims to understand:

- How lion tamarins use the cabruças in space and time
- Which types of cabruça supports a viable population.
- Which parameter of the maintenance is crucial for the suitability as habitat.



A lion tamarin of Fazenda Bom Pastor equipped with a radio collar.

The project is a collaboration between AMAP Brazil and the Universidade Estadual de Santa Cruz (UESC). Dr. Teixeira coordinates this project as project coordinator of AMAP Brazil and Post Doc at the UESC. The project is carried out by research assistants, employed by AMAP. Prof. Oliveira (UESC), collaborator and initiator of the project, holds all necessary permits to conduct the project. He is a member of the Primate

Specialist Group of IUCN. The results will be published and incorporated into the [IUCN](#) assessment of the conservation status of *L. chrysomelas* and in the Ministry of Environment's National [Action Plans](#) for Mammal and Primate Conservation.

The results will allow AMAP to create a guideline for local cocoa farmers to practice biodiversity-friendly cocoa harvesting and maintenance.



Jiomario "Bila" dos Santos Souza with TR-4K receiver and handheld antenna.

Target groups

The project is directed at the following audiences:

Local level:

Local stakeholders, mainly cocoa farmers and landowners whose land is or could be a habitat of golden-headed lion tamarins.

National level:

Administrative actors, such as ICMBio (Institute for Biodiversity Conservation), which develops the national conservation plans.

International level:

The research community working on the conservation of Brazil's primates, the IUCN - to assess the conservation status of golden-headed lion tamarins.

II. Project Design

2.1 Study area

The study area in the municipality Ilhéus is dominated by cabruca as landscape matrix, with patches of secondary forest, pastures, and is situated on Fazenda Bom Pastor and the

neighboring cocoa-farms Santa Rita and Bom Fin/Almada. These farms are actively managed cocoa farms with varying proportions of cabruca, secondary forest and pasture. Several habituated lion tamarin groups are monitored, named after the Fazendas where they are most frequently found.



Fazenda Bom Pastor, monitoring project area in the original distribution area of golden-headed lion tamarins in Bahia, Brazil

2.2 Methods

Radio Telemetry is used to facilitate location and monitoring. On one or two individuals of each lion tamarin groups, have attached RI-2D radio transmitter collars (Holohil Inc.), optimized for small mammals with a body weight of 550-590gr. The capture for the collar attachment is carried out by using Tomahawk live traps (48.3 x 15.2 x 15.2 cm), designed for small mammals, baited with bananas and placed on wooden platforms 1,5m above the ground. The capture takes place once or twice a year and is carried out only by trained personal under the supervision of a veterinarian (see paragraph III). Transmitter collared lion tamaris are detected in its habitat by using a biomedical receiver TR-4K (Telonics Inc.) and a handheld antenna.

2.3 Data Collection

During the capture for each individual following data are recorded:

Weight, knee to heel and wrist to elbow length, reproductive condition and group size and composition (age and sex). The group size and composition is adjusted by including not captured individuals in the surrounding of the traps. As cooperative breeders that lives in

family groups, it is assumed, that not captured individuals of the group remain in the vicinity of captured individuals. Tooth wear is used to estimate the age of adult animals and tooth wear, body weight and dental composition to estimate ages of younger members of group. The monitoring takes place once a week for each group. A group is monitored the whole day, starting when they left a sleeping site in the morning until they enter a sleeping site in the evening, or on partial days either from the morning until noon or from noon until evening.



Preparing a platform for catching the golden-headed lion tamarins

Following data are documented:

Demographic data:

- Number of Individuals (male, female, offspring, infants)
- Reproduction (time of breeding, number and time of offspring)

Ecological data:

- Diets
- Sleeping site use
- Home range

Behavioral data:

- Amount of time spend with foraging in bromeliads, fruit trees and others
- Amount of time spend with resting
- Moving distance
- Moving speed
- Changing of sleeping sites

III. The project team

The project is coordinated by Dr. Joanison Vicente dos Santos Teixeira and carried out by two research assistants. Students are integrated into the team as part of master's or doctoral theses.

Dr. Joanison Vicente dos Santos Teixeira

[Dr. Teixeira](#) holds an unpaid post doc position at LECAP (Laboratório de Etnoconservação e Áreas Protegidas) of the UESC. He is responsible for coordinating the project and evaluating and publishing the results. AMAP already supported the realization of his doctoral thesis. Since the beginning of 2022, Dr. Teixeira has been the Project Coordinator for golden-headed lion tamarin related projects for AMAP Brazil.

Prof. Leonardo de Carvalho Oliveira

Professor at the Universidade Estadual de Santa Cruz (UESC). He is member of the [Primate Specialist Group](#) of the IUCN. Prof. Oliveira began his work with golden-headed lion tamarins in 2008, and since then this primates have been the focus of his university work. Prof. Oliveira supervised also the PhD thesis of the project coordinator.

Prof. Danilo Simonini Teixeira

Institute of Veterinary Medicine at the UESC. [Prof. Simonini Teixeira](#) is a veterinarian and primatologist. His research group is responsible for the capture and local anaesthesia of the golden-headed lion tamarins for changing the collar transmitters. The state of health is also documented.

Students

Maria Alejandra, a student of the UESC will carry out her **master thesis** within the monitoring project. She will collect data on the ecology and behavior of the lion tamarins and will especially investigate the use of sleeping sites by lion tamarins. The data collection will begin in April 2024. The master thesis will take place as part of the regular monitoring at the fazendas Bom Pastor and Santa Rita. The study will use existing monitoring data, expand the data and analyses the dataset.

Research Assistants

Jiomario „Bila“ dos Santos Souza

Bila was hired by AMAP for this project in 2018. For over 20 years, Bila has been carrying out monitoring and guided tours on request of various scientists, including for the research work of [Becky Raboy](#) and [Kristel de Vleeschouwer](#). Since 2008, Bila has been working on behalf of Dr. Oliveira and is responsible for the realization of the monitoring in the field.

Rodrigo Souza dos Santos

was hired by AMAP as an assistant for Bila at the beginning of 2022 and was trained over the course of the year. He has already been working for AMAP in the reforestations since 2018, but showed a strong interest in working with golden-headed lion tamarins.

Their scope of work includes:

1. Conducting the basic monitoring
2. Catching the groups for the purpose of radio collar attachment and/or changing
3. Handling and maintaining of all equipment
4. Guidance of students under supervision of Prof. Oliveira and/or Dr. Teixeira.

VI. Animal handling methods

The lion tamarin groups will be captured and collared once or twice a year. Prof. Oliveira, has passed the ethics committee requirements to conduct safe captures and ensure the well-being of all individuals. Prof. Oliveira holds all necessary permits to conduct the monitoring with lion tamarins. Additionally, qualified veterinarians will accompany the capture process for all groups. All captures will be performed as described in Kleiman et al. 1986 and Dietz & Baker 1993, using non-injurious techniques which we are using since 2008.



Almada Mata Atlântica Projeto (AMAP)
Hofstatt 5, 35037 Marburg
amap-brazil.org

Our Partners



UMWELTSTIFTUNG | GREENPEACE



Wilhelma
Zoologisch-Botanischer Garten
Stuttgart



Artenschutzstiftung
Zoo Karlsruhe

V Publications generated with the projects monitoring data

- Almeida-Rocha de Monteiro, J., Pedreira dos Reis, P., de Carvalho Oliveira, L., 2014. Play Behavior of the Golden-Headed Lion Tamarin in Brazilian Cocoa Agroforests. *Folia Primatol* 85, 192–199. <https://doi.org/10.1159/000362813>
- Almeida-Rocha, J.M. de, Reis, P.P., de V. Grelle, C.E., Oliveira, L.C., 2015. Do Habitat Use and Interspecific Association Reflect Predation Risk for the Golden-Headed Lion Tamarin (*Leontopithecus chrysomelas*)? *Int J Primatol* 36, 1198–1215. <https://doi.org/10.1007/s10764-015-9885-6>
- Almeida-Rocha de Monteiro, Peres, C., Monsalvo, J., Oliveira, L., 2020. Habitat determinants of golden-headed lion tamarin (*Leontopithecus chrysomelas*) occupancy of cocoa agroforests: gloomy prospects for management intensification. *Am J Primatol*, in press
- Catenacci, L.S., Ferreira, M., Martins, L.C., De Vleeschouwer, K.M., Cassano, C.R., Oliveira, L.C., Canale, G., Deem, S.L., Tello, J.S., Parker, P., Vasconcelos, P.F.C., Travassos da Rosa, E.S., 2018. Surveillance of Arboviruses in Primates and Sloths in the Atlantic Forest, Bahia, Brazil. *EcoHealth* 15, 777–791. <https://doi.org/10.1007/s10393-018-1361-2>
- Costa, T. ; Nogueira-Filho, S. ; Vleeschouwer, K. ; Oliveira, Lc. ; Sousa, M. B. ; Mendl, M. ; Catenacci, L. S. ; Nogueira, S. S. C. 2020. Individual behavioral differences and health of golden headed lion tamarins. *American Journal of Primatology*, v. 82, p. 1-10, 2020.
- Magro Moraes, A., Grativol, A., D., De Vleeschouwer, K., M., Ruiz-Miranda, C., R., Raboy, B., E., Oliveira, L. , Dietz, M., Galbusera P., H., A., 2018. Population Genetic Structure of an Endangered Endemic Primate (*Leontopithecus chrysomelas*) in a Highly Fragmented Atlantic Coastal Rain Forest. *Folia Primatol* 89, 365–381. <https://doi.org/10.1159/000492176>
- Oliveira, L.C., Hankerson, S.J., Dietz, J.M., Raboy, B.E., 2010. Key tree species for the golden-headed lion tamarin and implications for shade-cocoa management in southern Bahia, Brazil. *Animal Conservation* 13, 60–70. <https://doi.org/10.1111/j.1469-1795.2009.00296.x>
- Oliveira, L.C., Dietz, J.M., 2011. Predation risk and the interspecific association of two Brazilian Atlantic forest primates in Cabruca agroforest. *Am. J. Primatol.* 73, 852–860. <https://doi.org/10.1002/ajp.20952>
- Oliveira, L.C., G. Neves, L., E. Raboy, B., M. Dietz, J., 2011. Abundance of Jackfruit (*Artocarpus heterophyllus*) Affects Group Characteristics and Use of Space by Golden-Headed Lion Tamarins (*Leontopithecus chrysomelas*) in Cabruca Agroforest. *Environmental Management* 48, 248–262. <https://doi.org/10.1007/s00267-010-9582-3>
- Oliveira, L.C., Estrada, A., 2017. Agroecosystems, in: Bezanson, M., MacKinnon, K.C., Riley, E., Campbell, C.J., Nekaris, K.A.I.A., Estrada, A., Di Fiore, A.F., Ross, S., Jones-Engel, L.E., Thierry, B., Sussman, R.W., Sanz, C., Loudon, J., Elton, S., Fuentes, A. (Eds.), *The International Encyclopedia of Primatology*. John Wiley & Sons, Inc., Hoboken, NJ, USA, pp. 1–6. <https://doi.org/10.1002/9781119179313.wbprim0235>
- Teixeira, Joanison Vicente Dos Santos, Fernando César Gonçalves Bonfim, Maurício Humberto Vancine, Milton C. Ribeiro, und Leonardo De C. Oliveira, 2023. „Effect of Landscape Attributes on the Occurrence of the Endangered Golden-headed Lion Tamarin in Southern Bahia, Brazil“. *American Journal of Primatology*, 25. Dezember 2023, e23588. <https://doi.org/10.1002/ajp.23588>.

Teixeira, Joanison Vicente Dos Santos, "Limites de distribuição, estimativa populacional e influência dos atributos da paisagem na ocorrência do ameaçado mico-leão-da-cara-dourada *Leontopithecus chrysomelas* (Kuhl, 1820), Primates; Callitrichidae", Doctoral Thesis, Universidade Estadual de Santa Cruz, 2022.

VI References

- Delabie, J.H.C., Jahyny, B., do Nascimento, I.C., Mariano, C.S.F., Lacau, S., Campiolo, S., Philpott, S.M., Leponce, M., 2007. Contribution of cocoa plantations to the conservation of native ants (Insecta: Hymenoptera: Formicidae) with a special emphasis on the Atlantic Forest fauna of southern Bahia, Brazil. *Biodivers Conserv* 16, 2359–2384. <https://doi.org/10.1007/s10531-007-9190-6>
- Dietz JM, Baker AJ (1993). Polygyny and female reproductive success in golden lion tamarins (*Leontopithecus rosalia*). *Animal Behaviour* 46: 1067-1078.
- Dietz JM, de Souza SN, Billerbeck R., 1996 .Population dynamics of golden-headed Lion tamarins *Leontopithecus chrysomelas* in Una Biological Reserve, Brazil. *Dodo, Journal of Wildlife Preservation Trust* 32:115–122
- Faria, D., Laps, R.R., Baumgarten, J., Cetra, M., 2006. Bat and Bird Assemblages from Forests and Shade Cacao Plantations in Two Contrasting Landscapes in the Atlantic Forest of Southern Bahia, Brazil. *Biodivers Conserv* 15, 587–612. <https://doi.org/10.1007/s10531-005-2089-1>
- Faria, D., Paciencia, M.L.B., Dixo, M., Laps, R.R., Baumgarten, J., 2007. Ferns, frogs, lizards, birds and bats in forest fragments and shade cacao plantations in two contrasting landscapes in the Atlantic forest, Brazil. *Biodivers Conserv* 16, 2335–2357. <https://doi.org/10.1007/s10531-007-9189-z>
- Holst, B., Medici, E.P., Marini-Filho, O.J., Kleiman, D., Leus, K., Pissinatti, A., Vivekananda, G., Ballou, J.D., Traylor-Holzer, K., Raboy, B., Passos, F., Vleeschouwer, K. & Montenegro, M.M. (Eds)., 2006. Lion tamarin population and habitat viability assessment workshop 2005. Final Report. Apple Valley, MN: IUCN/SSC Conservation Breeding Specialist Group.
- IUCN. The IUCN Red List of Threatened Species. Version 2016-3. <<http://www.iucnredlist.org>>. 2020.
- Kleiman, D.G., Beck, B.B., Dietz, J.M., Dietz, L.A., Ballou, J.D., Coimbra-Filho, A.F., 1986. Conservation Program for the Golden Lion Tamarin: Captive Research and Management, Ecological Studies, Educational Strategies, and Reintroduction, in: Benirschke, K. (Ed.), *Primates*. Springer New York, New York, NY, pp. 959–979. https://doi.org/10.1007/978-1-4612-4918-4_65
- Myers, N., Mittermeier, R.A., Mittermeier, C.G., da Fonseca, G.A.B., Kent, J., 2000. Biodiversity hotspots for conservation priorities. *Nature* 403, 853–858. <https://doi.org/10.1038/35002501>
- Oliveira, L.C., Dietz, J.M., 2011. Predation risk and the interspecific association of two Brazilian Atlantic forest primates in Cabruca agroforest. *Am. J. Primatol.* 73, 852–860. <https://doi.org/10.1002/ajp.20952>

- Oliveira, L.C., Estrada, A., 2017. Agroecosystems, in: Bezanson, M., MacKinnon, K.C., Riley, E., Campbell, C.J., Nekaris, K.A.I.A., Estrada, A., Di Fiore, A.F., Ross, S., Jones-Engel, L.E., Thierry, B., Sussman, R.W., Sanz, C., Loudon, J., Elton, S., Fuentes, A. (Eds.), *The International Encyclopedia of Primatology*. John Wiley & Sons, Inc., Hoboken, NJ, USA, pp. 1–6.
<https://doi.org/10.1002/9781119179313.wbprim0235>
- Oliveira, L.C., G. Neves, L., E. Raboy, B., M. Dietz, J., 2011. Abundance of Jackfruit (*Artocarpus heterophyllus*) Affects Group Characteristics and Use of Space by Golden-Headed Lion Tamarins (*Leontopithecus chrysomelas*) in Cabruca Agroforest. *Environmental Management* 48, 248–262. <https://doi.org/10.1007/s00267-010-9582-3>
- Oliveira, L.C., Hankerson, S.J., Dietz, J.M., Raboy, B.E., 2010. Key tree species for the golden-headed lion tamarin and implications for shade-cocoa management in southern Bahia, Brazil. *Animal Conservation* 13, 60–70. <https://doi.org/10.1111/j.1469-1795.2009.00296.x>
- Pardini, R., 2004. Effects of forest fragmentation on small mammals in an Atlantic Forest landscape. *Biodiversity and Conservation* 13, 2567–2586.
<https://doi.org/10.1023/B:BIOC.0000048452.18878.2d>
- Pinto, L.P. de S., Rylands, A.B., 1997. Geographic Distribution of the Golden-Headed Lion Tamarin, *Leontopithecus chrysomelas*: Implications for Its Management and Conservation. *Folia Primatol* 68, 161–180. <https://doi.org/10.1159/000157244>
- Rylands, A.B., 1993. The ecology of the lion tamarins, *Leontopithecus*: some intrageneric differences and comparisons with other callitrichids. In *Marmosets and tamarins: systematics, behaviour and ecology*: 296–313. Rylands, A.B. (Ed.). Oxford: Oxford University Press.
- Sambuichi R.H.R., 2002. Fitossociologia e diversidade de espécies arbóreas em cabruca (Mata Atlântica raleada sobre plantação de cacau) na região sul da Bahia, Brasil. *Acta Botanica Brasiliensis* 16:89–101
- Sambuichi R.H.R., 2006. Estrutura e dinâmica do componente arbóreo em área de cabruca na região cacauífera do sul da Bahia, Brasil. *Acta Botanica Brasiliensis* 20:943–954
- Sambuichi, R.H.R., Haridasan, M., 2007. Recovery of species richness and conservation of native Atlantic forest trees in the cacao plantations of southern Bahia in Brazil. *Biodivers Conserv* 16, 3681–3701. <https://doi.org/10.1007/s10531-006-9017-x>
- Schroth, G., do Socorro Souza da Mota, M., 2014. Agroforestry: Complex Multistrata Agriculture, in: *Encyclopedia of Agriculture and Food Systems*. Elsevier, pp. 195–207.
<https://doi.org/10.1016/B978-0-444-52512-3.00030-9>
- Schroth, G., Harvey, C.A., 2007. Biodiversity conservation in cocoa production landscapes: an overview. *Biodivers Conserv* 16, 2237–2244. <https://doi.org/10.1007/s10531-007-9195-1>
- Shi, H., Singh, A., Kant, S., Zhu, Z., Waller, E., 2005. Integrating Habitat Status, Human Population Pressure, and Protection Status into Biodiversity Conservation Priority Setting: *Integrating Social Factors into Priority Setting*. *Conservation Biology* 19, 1273–1285.
<https://doi.org/10.1111/j.1523-1739.2005.00225.x>
- Zeigler, S.L., Fagan, W.F., DeFries, R., Raboy, B.E., 2010. Identifying Important Forest Patches for the Long-Term Persistence of the Endangered Golden-Headed Lion Tamarin (*Leontopithecus chrysomelas*). *Tropical Conservation Science* 3, 63–77.
<https://doi.org/10.1177/194008291000300106>